

**Table 1**

Littering Parameters for Wistar Han rats after in utero/postnatal exposure.

| Dose (mg/kg)  | 0                     | 3        | 15       | 50       |
|---|-----------------------|----------|----------|----------|
| Time-Mated Females (GD 6)                             | 62                    | 52       | 52       | 62       |
| Females Pregnant (%)                                  | 54 (87%)              | 42 (81%) | 43 (83%) | 51 (82%) |
| Dams with Evidence of Pregnancy<br>Not Delivering (%) | 2 (4%)                | 1 (2%)   | 4 (9%)   | 2 (4%)   |
| Dams with Litters on PND 0 (%)                        | 52 (96%)              | 41 (98%) | 39 (91%) | 49 (96%) |
| Dams, Moribund or Natural<br>Deaths                   | 0                     | 0        | 0        | 0        |
| Litters Post-Standardization (PND<br>4)               | 36                    | 29       | 28       | 37       |
| Post-Weaning Allocation                               |                       |          |          |          |
| F1 Males – Core (litters)                             | 50 (29 <sup>a</sup> ) | 50 (25)  | 50 (25)  | 50 (29)  |
| F1 Females – Core (litters)                           | 50 (30)               | 50 (25)  | 50 (25)  | 50 (28)  |

There were no significant differences among the groups for any of the litter parameters.

<sup>a</sup> Number of litters from which pups were taken at PND22 to populated the 50 rats/sex/dose group that went on for 2 year studies.