

Table 6
Incidence of Cardiac Lesions at PND 21.

AGE	Sex/Pathology	0	BPA 2.5	BPA 25	BPA 250	BPA 2500	BPA 25000	EE2 0.05	EE2 0.5
PND 21	Male								
	Cardiomyopathy	9/10 (1.3)	8/8 (1.1)	8/8 (1.1)	8/9 (1.2)	8/10 (1.2)	8/10 (1.3)	10/10 (1.5)	10/10 (1.7)
	Focal Fibrosis	6/10	4/8	3/8	2/9	3/10	6/10	6/10	3/10
	Inflammation/Necrosis	3/10	3/8	3/8	2/9	4/10	3/10	4/10	4/10
	Diffuse Degeration	–	–	–	1/9	–	–	–	–
	Female								
	Cardiomyopathy	6/10 (0.8)	10/10 (1.8) ^a	9/10 (1.4)	10/10 (1.8) ^a	7/8 (1.1)	10/10 (1.9) ^b	10/10 (2.3) ^c	10/10 (2.0) ^d
	Focal Fibrosis	5/10	7/10	6/10	4/10	3/8	4/10	4/10	6/10
	Inflammation/Necrosis	2/10	6/10	5/10	3/10	3/8	5/10	6/10	3/10
	Diffuse Degeration	–	1/10	1/10	–	–	–	–	–

Shown in parentheses are the mean severity scores. Severity scores for control and each exposure group were examined using the rank order ANOVA Kruskal-Wallis H test with Dunn's multiple comparisons tests.

^a p = .0536, d = 1.34.

^b p = .02, d = 1.52.

^c p = .0005, d = 2.42.

^d p = .01, d = 1.73.