

**Table S2: Number of days (d) spent in estrus, diestrus and proestrus following in utero exposure to bisphenol A (BPA), bisphenol AF (BPAF), or bisphenol S (BPS) during postnatal days (PND) 63-83**

	Estrus (d)	Diestrus (d)	Proestrus (d)
Vehicle	7.2 ± 4.0	11.7 ± 3.3	2.0 ± 1.0
BPA 0.5 mg/kg	7.6 ± 2.9	11.4 ± 2.7	1.2 ± 1.3
BPA 5 mg/kg	7.7 ± 2.7	11.0 ± 2.4	1.0 ± 0.6*
BPA 50 mg/kg	6.1 ± 2.4	12.1 ± 3.1	1.4 ± 1.2
BPAF 0.05 mg/kg	7.3 ± 2.2	11.2 ± 2.9	1.5 ± 1.2
BPAF 0.5 mg/kg	7.0 ± 2.3	11.7 ± 2.6	1.6 ± 1.0
BPAF 5 mg/kg	5.6 ± 2.4	12.8 ± 1.4	1.4 ± 0.7
BPS 0.05 mg/kg	7.5 ± 3.3	11.2 ± 3.8	1.7 ± 1.0
BPS 0.5 mg/kg	6.0 ± 3.7	12.4 ± 4.0	1.3 ± 1.3
BPS 5 mg/kg	6.7 ± 2.0	11.4 ± 2.2	1.7 ± 1.3

Data are presented as the mean number of days in that stage ± SEM.

Litter numbers: Vehicle (n = 9), BPA 0.5 (n = 12), BPA 5 (n = 12), BPA 50 (n = 9), BPAF 0.05 (n = 9),