

Study Number: MOG10866
Test Type: MOG - Range Finding
Route: Dosing in Feed
Species/Strain: Rat/Harlan Sprague Dawley

R16: Pubertal Markers Summary
Test Compound: Isopropylated Phenyl Phosphate
CAS Number: 68937-41-7

Date Report Requested: 02/19/2021
Time Report Requested: 13:49:25
Lab: Battelle

Study Number: MOG10866
Study Gender: Both
PWG Approval Date: See web page for date of PWG Approval
Version: v1.1.7

Study Number: MOG10866

Test Type: MOG - Range Finding

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

R16: Pubertal Markers Summary

Test Compound: Isopropylated Phenyl Phosphate

CAS Number: 68937-41-7

Date Report Requested: 02/19/2021

Time Report Requested: 13:49:25

Lab: Battelle

		Male			
Generation	Cohort	Treatment Groups (ppm)			
		0	1000	3000	
F1	All Males	No. Examined (litters)	32 (7)	60 (16)	17 (7)
		No. Removed (litters)	0 (0)	0 (0)	0 (0)
		No. Not Attaining BPS (litters)	0 (0)	0 (0)	0 (0)
		Day of BPS			
		Mean Analysis			
		Litter Mean \pm SE	41.6 \pm 0.6 *	42.0 \pm 0.4	43.1 \pm 0.7
		Litter Mean of Adjusted \pm SE	41.8 \pm 0.5	42.2 \pm 0.4	43.0 \pm 0.7
		Proportional Hazards Analysis			
		Litter-based Model	p=0.014	p=0.495	p=0.172
		BW at Attainment (g)	167.2 \pm 3.8 *	164.6 \pm 3.9	151.6 \pm 5.1
BW at Weaning (g)	77.1 \pm 3.3	77.2 \pm 2.5	70.8 \pm 3.3		

Study Number: MOG10866

Test Type: MOG - Range Finding

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

R16: Pubertal Markers Summary

Test Compound: Isopropylated Phenyl Phosphate

CAS Number: 68937-41-7

Date Report Requested: 02/19/2021

Time Report Requested: 13:49:25

Lab: Battelle

		Female			
Generation	Cohort	Treatment Groups (ppm)			
		0	1000	3000	
F1	All Females	No. Examined (litters)	38 (10)	43 (14)	23 (7)
		No. Removed (litters)	33 (10)	28 (10)	15 (8)
		No. Not Attaining VO (litters)	0 (0)	0 (0)	0 (0)
		Day of VO			
		Mean Analysis			
		Litter Mean ± SE	36.4 ± 0.5 **	36.5 ± 0.4	38.8 ± 0.7 *
		Litter Mean of Adjusted ± SE	36.6 ± 0.4 **	36.6 ± 0.4	38.8 ± 0.7 *
		Proportional Hazards Analysis			
		Litter-based Model	p=0.002	p=0.422	p=0.022
		BW at Attainment (g)	111.2 ± 2.5	108.0 ± 2.8	115.0 ± 4.4
BW at Weaning (g)	70.4 ± 3.0	68.9 ± 2.1	67.2 ± 3.0		

Study Number: MOG10866

Test Type: MOG - Range Finding

Route: Dosing in Feed

Species/Strain: Rat/Harlan Sprague Dawley

R16: Pubertal Markers Summary

Test Compound: Isopropylated Phenyl Phosphate

CAS Number: 68937-41-7

Date Report Requested: 02/19/2021

Time Report Requested: 13:49:25

Lab: Battelle

LEGEND

BPS = Balanopreputial separation; BW = Body weight; VO = Vaginal opening

No. Examined (litters) = the number of animals or pups examined (number of litters)

No. Removed (litters) is the number of animals (number of litters contributing) that died or were removed prior to the end of the observation period and did not attain. These animals were excluded from all analyses.

No. Not Attaining BPS (litters) and No. Not Attaining VO (litters) is the number of animals (number of litters contributing) that survived to the end of the observation period without attaining.

Summary statistics and mixed model results are presented for animals that attained during the observation period for Day of BPS and Day of VO Mean Analysis endpoint.

Means of litter means presented for Day of BPS and Day of VO Litter Mean \pm SE. Trend and pairwise tests were based on mixed models for day of attainment with dose as a covariate and a random effect for litter. The Dunnett-Hsu adjustment was used for multiple comparisons.

Mean adjusted day of attainment was calculated from the mean of the litter means of the weaning weight-adjusted attainment days for individual pups. Trend and pairwise tests were based on mixed models for day of attainment with dose and weaning weight as covariates and a random effect for litter. The Dunnett-Hsu adjustment was used for multiple comparisons.

Animals that did not attain by the end of the observation period were included in the proportional hazards analysis.

P-values for trend and pairwise comparisons for the Litter-based Model of the Proportional Hazards Analysis were calculated from a Cox proportional hazards model with dose and weaning weight as covariates and a random effect for litter, and a Hommel adjustment for multiple comparisons.

Analysis of body weight at attainment and body weight at weaning were performed using mixed effects models with dose as covariate and a random effect for litter. The Dunnett-Hsu adjustment was used for multiple comparisons. Animals that attained during the observation period were used for analysis.

Statistical significance for the control group indicates a significant trend test

Statistical significance for a treatment group indicates a significant pairwise test compared to the vehicle control group

* Statistically significant at $P \leq 0.05$

** Statistically significant at $P \leq 0.01$

10000 ppm VO data was excluded from analysis.

The 30,000 ppm group was terminated on GD12, and the 15,000 ppm group was terminated by LD3/PND3 due to excessive toxicity.

**** END OF REPORT ****