

**Table 1.** Data sources

<b>Test method</b>	<b>Reference</b>
DPRA	<p>Bauch <i>et al.</i> (2011)</p> <p>Bauch <i>et al.</i> (2012)</p> <p>Gerberick <i>et al.</i> (2004)</p> <p>Gerberick <i>et al.</i> (2007)</p> <p>Jaworska <i>et al.</i> (2011)</p> <p>Jaworska <i>et al.</i> (2013)</p> <p>Joint Research Centre of the European Union (2013)</p> <p>Natsch <i>et al.</i> (2013)</p> <p>Nukada <i>et al.</i> (2013)</p>
KeratinoSens	<p>Ball <i>et al.</i> (2011)</p> <p>Bauch <i>et al.</i> (2011)</p> <p>Bauch <i>et al.</i> (2012)</p> <p>Emter <i>et al.</i> (2010)</p> <p>Joint Research Centre of the European Union (2014)</p> <p>Natsch <i>et al.</i> (2013)</p>
h-CLAT	<p>Ashikaga <i>et al.</i> (2010)</p> <p>Bauch <i>et al.</i> (2011)</p> <p>Bauch <i>et al.</i> (2012)</p> <p>Nukada <i>et al.</i> (2011)</p> <p>Nukada <i>et al.</i> (2012)</p> <p>Nukada <i>et al.</i> (2013)</p> <p>Sakaguchi <i>et al.</i> (2010)</p> <p>Takenouchi <i>et al.</i> (2013)</p>

<b>Test method</b>	<b>Reference</b>
LLNA	Basketter <i>et al.</i> (1996) and Estrada <i>et al.</i> (2003) (xylene)  Basketter and Kimber (2006) (diphenylcyclopropanone, maleic anhydride, and propyl gallate)  NICEATM LLNA database  Van Och <i>et al.</i> (2000) (phthalic anhydride)

---

DPRA, direct peptide reactivity assay; h-CLAT, human cell line activation test; LLNA, murine local lymph node assay; NICEATM, National Toxicology Program Interagency Center for the Evaluation of Alternative Toxicological Methods.