

**Table 1.** Selected Organ Weight and Organ Weight to Body Weight Ratios in Mice in the 90-d Gavage Study of CYN<sup>a</sup>.

Dose ( $\mu\text{g}/\text{kg}$ )	0	75	150	300
<b>Male</b>				
<i>n</i>	10	10	9	10
<b>Necropsy weight</b>	40.6 $\pm$ 1.4	40.4 $\pm$ 0.9	38.1 $\pm$ 0.6	39.3 $\pm$ 0.7
<b>Liver</b>				
Absolute	1.96 $\pm$ 0.07	2.37 $\pm$ 0.08**	2.36 $\pm$ 0.05*	2.70 $\pm$ 0.06*
Relative	0.048 $\pm$ 0.002	0.059 $\pm$ 0.001*	0.062 $\pm$ 0.001*	0.069 $\pm$ 0.001*
<b>Kidney</b>				
Absolute	0.62 $\pm$ 0.03	0.88 $\pm$ 0.03*	0.89 $\pm$ 0.04*	0.84 $\pm$ 0.05*
Relative	0.015 $\pm$ 0.001	0.022 $\pm$ 0.001*	0.023 $\pm$ 0.001*	0.022 $\pm$ 0.001*
<b>Testes</b>				
Absolute	0.248 $\pm$ 0.009	0.246 $\pm$ 0.011	0.252 $\pm$ 0.010	0.277 $\pm$ 0.013*
Relative	0.0061 $\pm$ 0.0002	0.0061 $\pm$ 0.0002	0.0067 $\pm$ 0.0002	0.0071 $\pm$ 0.0004*
<b>Female</b>				
<i>n</i>	9	10	10	10
<b>Necropsy weight</b>	33.5 $\pm$ 1.7	31.8 $\pm$ 1.1	30.5 $\pm$ 0.6	29.8 $\pm$ 0.7*
<b>Liver</b>				
Absolute	1.64 $\pm$ 0.11	1.66 $\pm$ 0.07	1.88 $\pm$ 0.09*	1.87 $\pm$ 0.05*
Relative	0.049 $\pm$ 0.001	0.052 $\pm$ 0.001	0.062 $\pm$ 0.003*	0.063 $\pm$ 0.001*
<b>Kidney</b>				
Absolute	0.38 $\pm$ 0.015	0.39 $\pm$ 0.01	0.40 $\pm$ 0.02	0.39 $\pm$ 0.01
Relative	0.012 $\pm$ 0.000	0.012 $\pm$ 0.000	0.013 $\pm$ 0.001*	0.013 $\pm$ 0.000*

<sup>a</sup>Necropsy body weight and organ weights are given in grams (g); relative organ weights (organ weight to body weight ratios) are given as organ weight (g)/body weight (g) (mean  $\pm$  standard error).

\*Significant differences ( $p \leq .05$ ) from control using Wilcoxon and Kruskal–Wallis tests.

\*\*Significant differences ( $p \leq 0.01$ ) from control using Wilcoxon and Kruskal–Wallis tests.